Supplemental Online Content

Demer AM, Hanson JL, Maher IA, Liszewski W. Association of Mohs micrographic surgery vs wide local excision with overall survival outcomes for patients with melanoma of the trunk and extremities. *JAMA Dermatol*. Published online October 21, 2020. doi:10.1001/jamadermatol.2020.3950

eTable. Baseline Treatment Group Characteristics

This supplemental material has been provided by the authors to give readers additional information about their work.

Supplemental Materials

eTable. Baseline Treatment Group Characteristics

| e rabie. Daseille | Treatment Group Characteristics | | | |
|----------------------|---------------------------------|-----------------|---------|--|
| | MMS | WLE | p-value | |
| Number treated | 4,413 (2.3%) | 184,449 (97.7%) | | |
| Age (Mean (SD)) | 59.3 (15.9) | 58.7 (16.0) | 0.037 | |
| Sex | | | | |
| Male | 2,239 (50.7%) | 97,209 (52.7%) | <0.010 | |
| Female | 2,174 (49.3%) | 87,204 (47.3%) | | |
| Race | | | | |
| White | 4,292 (97.3%) | 180,038 (97.6%) | 0.133 | |
| Other | 121 (2.7%) | 4,411 (2.4%) | | |
| Insurance | | | | |
| Private | 2,566 (58.1%) | 109,768 59.5%) | 0.027 | |
| None | 102 (2.3%) | 4,602 (2.5%) | | |
| Medicaid | 88 (2.0%) | 4,204 (2.3%) | | |
| Medicare | 1,609 (36.5%) | 64,087 (34.7%) | | |
| Other gov. insurance | 48 (1.1%) | 1,788 (1.0%) | | |
| Location | | | | |
| Trunk | 1,767 (40.0%) | 77,547 (42.0%) | 0.318 | |
| Upper Extremity | 1,595 (36.1%) | 61,520 (33.4%) | | |
| Lower extremity | 1,051 (23.8%) | 45,382 (24.6%) | | |
| Margins | | | | |
| Negative | 4,315 (97.8%) | 180,950 (98.1%) | 0.120 | |
| Positive | 98 (2.2%) | 3,499 (1.9%) | | |
| Ulceration | | | | |
| No | 3,940 (89.3%) | 152,120 (82.5%) | <0.001 | |
| Yes | 473 (10.7%) | 32,329 (17.5%) | | |
| Charlson-Deyo | | | | |
| 0 | 3,961 (89.8%) | 162,378 (88.0%) | 0.002 | |
| 1 | 372 (8.4%) | 18,414 (10.0%) | | |
| 2 or more | 80 (1.8%) | 3,657 (2.0%) | | |
| Tumor type | | | | |
| Nodular | 221 (5.0%) | 18,555 (10.1%) | <0.001 | |
| Other | 2,330 (52.8%) | 94,043 (51.0%) | | |
| | • | • | - | |

| LM/LMM | 286 (6.5%) | 5,189 (2.8%) | |
|------------------------------|---------------|----------------|--------|
| Superficial Spreading | 1,576 (35.7%) | 66,662 (36.1%) | |
| Breslow Depth (Mean (SD)) | 0.91 (1.45) | 1.42 (1.83) | <0.001 |

Statistically significant differences in age, sex, insurance status, ulceration, Charlson-Deyo comorbidity score, tumor type, and Breslow depth were identified between treatment groups. Importantly, each of these characteristics were controlled for on multivariate analysis. No difference in tumor location or race were identified between MMS and WLE cohorts.